

Andrew County Health Department
Point of Distribution (POD)

POD Location

Andrew County Health Departments POD is a “Drive Through”. It will be at the South entrance to the High School off State Route E.

For more information listen to and watch the local television and radio stations for public service announcements.

If a **Point of Distribution (POD)** is opened do to any emergency where medications are dispensed, print this form out. Fill it out and bring it with you to the POD.

NOTE: Only one adult from the house hold needs to bring the form.

1		2 FOR EACH MEMBER OF YOUR HOUSEHOLD, ANSWER ALL THREE QUESTIONS BELOW:						
		A	B	C	DECISION MATRIX – STAFF USE ONLY			
LIST ALL HOUSEHOLD MEMBERS FOR WHOM YOU ARE PICKING UP MEDICATIONS TODAY, INCLUDING YOURSELF		Is household member:	Is household member allergic to or shouldn't take any of these:	Is household member allergic to or shouldn't take any of these:	Answer A	Answer B	Answer C	Provide
		<ul style="list-style-type: none"> • Pregnant • Breast feeding • Under 8 years of age 	<ul style="list-style-type: none"> • Doxycycline (Vibramycin) • Minocycline • Tetracycline 	<ul style="list-style-type: none"> • Ciprofloxacin • Levofloxacin (Levaquin) • Ofloxacin • Gatifloxacin • Moxifloxacin 	No	No / DK	No / DK	Doxy
					Yes / DK	No / DK	No / DK	Cipro
					Yes / DK	No / DK	Yes	Doxy
					Yes / DK	Yes	No / DK	Cipro
					Yes / DK	Yes	Yes	Refer
					No	No / DK	Yes	Doxy
					No	Yes	No / DK	Cipro
					No	Yes	Yes	Refer
		Last name	First Name	Yes, No or Don't Know?	Yes, No or Don't Know?	Yes, No or Don't Know?	CIRCLE MEDICATION TO BE PROVIDED STAFF USE ONLY	
					Doxy	Cipro	Refer	
					Doxy	Cipro	Refer	
					Doxy	Cipro	Refer	
					Doxy	Cipro	Refer	
					Doxy	Cipro	Refer	
					Doxy	Cipro	Refer	
					Doxy	Cipro	Refer	
Add Totals Under Doxy & Cipro Columns:								
NCR Medication Screening Form		3 EACH PERSON SHOULD TAKE THE MEDICINE CIRCLED IN THEIR ROW.						

POD SPECIFIC INSTRUCTIONS

The FDA instructions for **Doxycycline** (which would provide the proper doses for anthrax, plague, and tularemia prophylaxis), are available on our SNS website at <http://www.dhss.mo.gov/StrategicNationalStockpile/15-TabletCrushingInstructions-Doxy.pdf>. The related FDA website is found at http://www.fda.gov/cder/drug/infopage/penG_doxy/home_prep.htm.

Instructions for **ciprofloxacin** are, with slight modifications, reproduced from documents prepared by the Oklahoma City-County Health Department, and are available on our SNS website. Because the dosages for post-exposure prophylaxis for anthrax, plague, and tularemia differ, there is a separate set of instructions for each of these diseases.

Ciprofloxacin used as post-exposure prophylaxis for anthrax:

<http://www.dhss.mo.gov/StrategicNationalStockpile/09-AnthraxTabletCrushingInstructions-Cipro.pdf>

Ciprofloxacin used as post-exposure prophylaxis for plague:

<http://www.dhss.mo.gov/StrategicNationalStockpile/26-CiproPedPlague.pdf>

Ciprofloxacin used as post-exposure prophylaxis for tularemia:

<http://www.dhss.mo.gov/StrategicNationalStockpile/27-CiproPedTularemia.pdf>